Getting started on VENCLYXTO[®] (venetoclax)



Information in this booklet is up to date at the time of going to print. For further information on this medicine please refer to the Package Leaflet that comes with your medication.



About this guide

Your doctor has prescribed VENCLYXTO® (venetoclax) as the next stage of your treatment plan for Chronic Lymphocytic Leukaemia (CLL). Your medicine may be given to you in combination with other medicines or alone.

In this guide you'll learn about how your medicine works, and how and when to take it. You'll also find out how to help minimise the risk of experiencing certain side effects, as well as how to manage them if they do occur.

There is a treatment plan to fill in during the first few weeks of treatment. It is an easy way to remember what dose of your medicine to take each week and any other special instructions that your healthcare team has for you. You can also use it to record any side effects you might experience and note down any questions you'd like to ask your healthcare team.



Remember, your healthcare team is your primary support every step of the way. They can help you with any questions or concerns about your treatment.



Hospital/Clinic:	 	
Address:	 	
Email:	 	
Consultant:	 	
Specialist nurse:	 	
Phone no:	 	
After hours phone no:	 	



GP's name:	 	 	_	 	 	 	 	-	 -	 -	 -	 	-	 	 	
Email:	 	 		 	 	 	 		 _	 -	 	 	_	 	 	
Phone no:	 	 		 	 	 	 		 	 	 	 		 	 	



In this guide

- 2 How VENCLYXTO® works to fight CLL
- 4 How to take and store your medicine
- 10 Managing possible side effects
- 18 Your treatment plan
- 49 Your patient information booklet



How VENCLYXTO[®] works to fight CLL

Your medicine is taken in tablet form, once a day. It is a type of medication known as a 'BCL2 inhibitor'.



Normal cells in your body contain proteins called 'BCL2'.



CLL cells make too much BCL2, which helps the cancer cells to survive.





VENCLYXTO® works by blocking (inhibiting) BCL2 in cancer cells that are producing too much of the protein. Blocking this protein helps to kill and lower the number of cancer cells. It also slows down the worsening of the disease.

Please speak to your healthcare team if you have any questions.

How to take and store your medicine

What tablet(s) do I take each day?

Your medicine is always taken once a day in tablet form.

The exact number of tablets you'll take per day will depend on your healthcare team. Your healthcare team will start you at a low dose for 1 week. Your healthcare team will gradually increase the dose over the next 4 weeks to the full standard dose. For the first 4 weeks you will get a new pack each week.

It is very important that you **follow your healthcare team's instructions for the gradual increase in dosage** as it has been designed to help reduce the chance of getting something known as 'tumour lysis syndrome' (TLS), which you'll learn more about in the next section.

Ĵ

The information on these pages is a guide only, and you should always take the medicine exactly as your healthcare team or pharmacist has told you. You can always check with them if you are unsure. This diagram shows what you can expect to take in the first 5 weeks of treatment. You'll get a new pack of tablets each week for the first 5 weeks.



Images are for illustrative purposes only.

After the 5th week, you may be on the 400mg (4 x 100mg tablets) daily dose for as long as your healthcare team thinks is necessary. Occasionally your healthcare team may tell you to take a lower dose for a short period of time, but in general you should take 400mg daily as the 'maintenance dose'.

How and when do I take VENCLYXTO®?

As well as following the dosage prescribed by your healthcare team, you should also keep some things in mind every time you take your medicine:



Your first dose may be given **in hospital** under the supervision of your healthcare team. They will check your blood test results after 6 to 8 hours and 24 hours to see whether they need to adjust the dose.

So, on the second day, remember to **wait until your healthcare team tells you how many tablets to take**. This might also be necessary with subsequent dose increases, but your healthcare team will let you know.

What other important things should I remember?



Drink plenty of water when taking your medicine during the first 5 weeks of treatment. You should start drinking at least 1.5 to 2 litres (L) of water daily, two days before starting this medicine.



Do not eat grapefruit products, Seville oranges (bitter oranges) or star fruit (carambola) while you are taking your medicine. This includes eating them, drinking their juice, or taking a supplement that might contain them. This is because they can increase the amount of VENCLYXTO[®] in your blood.



If you vomit after taking your medicine, do not take an extra dose that day. Take the next dose at the usual time the next day. If you have problems taking this medicine, talk to your doctor.



Do not take more your medicine than you should. If you do, talk to your healthcare team or pharmacist, or go to the hospital immediately. Take the tablets and the package leaflet with you.



Do not stop taking your medicine unless your doctor tells you to.



Important: If you are taking any other medication or start a new medicine while you are taking VENCLYXTO[®], tell your healthcare team.

How do I store VENCLYXTO®?



Your medicine does not require any special storage conditions.



Make sure that your medicine is stored out of the sight and reach of children.



Do not use your medicine after the expiry date. The expiry date is stated on the carton and blister pack after 'EXP'.



Do not throw away any medicines via wastewater or household waste.

Ask your pharmacist how to dispose of medicines you no longer use.



What should I do if I miss a dose?



If it is **less than 8 hours** since the time you usually take your dose, you should take the missed dose as soon as possible.



If it is **more than 8 hours** since the time you usually take your dose, **do not** take the missed dose. Instead, return to your normal dose schedule the next day. Let your healthcare team know that you have missed a dose.



Never take a double dose to make up for a **missed dose.** If you are unsure, talk to your healthcare team or pharmacist.

What should I do if I don't take enough of this medicine?

It's very important to take your tablets exactly as your healthcare team prescribes in order to get the most benefit out of your medicine. If you don't have enough tablets for a dose, **contact your healthcare team or pharmacist as soon as possible**.

Managing possible side effects

What side effects are associated with VENCLYXTO®?

Like all medicines, your medicine can cause side effects. Not everyone will get them but it's important to be aware of what they are so you can get help from your healthcare team right away if necessary.

Although some side effects are known to occur with this medicine, it is still important to tell your healthcare team if you experience any of them.



Remember: You have an important role to play in your treatment. Drink plenty of water. It helps to remove cancer cell breakdown products from your blood through your urine.

If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the Package Leaflet. You can also report side effects directly to the Health Products Regulatory Authority (HPRA) Pharmacovigilance department; website: www.HPRA.ie. By reporting side effects, you can help provide more information on the safety of medicines.

Common side effects

You may experience some very common side effects (may affect more than one in ten people). Remember to tell your healthcare team about any changes in your health, including:

- Pneumonia
- Any signs of an upper respiratory tract infection, such as a runny nose, sore throat or cough
- Diarrhoea
- Feeling sick or being sick (nausea or vomiting)
- Constipation
- Feeling tired

Other side effects

There are other side effects that occur commonly (may affect up to one in ten people). It is important to get help from your healthcare team if you get any of these:

- Severe infection in the blood (sepsis)
- Urinary tract infection
- Low number of white blood cells with fever (febrile neutropenia)

Blood test may also show:

- higher level of creatinine
- higher level of urea



At the first sign of infection or feeling unwell, contact your healthcare team.



Neutropenia

Low white blood cell count (neutropenia) and infections are very common and may effect more than 1 in 10 people.

You may also hear it called a low neutrophil count. It's important to attend all your appointments and to have all your blood tests taken on time, as these allow your healthcare team to check your white blood cell count routinely. It is important to do these checks because healthy white blood cells help your body to fight off infections.

Your healthcare team will check your blood count during treatment with your medicine. Low white blood cell count can increase your risk for infection. Signs may include fever, chills, feeling weak or confused, cough, pain or burning feeling when passing urine. Some infections can be serious and may lead to death. Tell your healthcare team immediately if you have signs of an infection while taking this medicine.

Tumour Lysis Syndrome (TLS)

Some people receiving treatment for CLL can develop tumour lysis syndrome (TLS), which is caused by the fast breakdown and destruction of cancer cells as a result of treatment.

Because of the way your medicine works, TLS and its side effects may occur in some patients (up to one in ten people). This is because, as the cancer cells are destroyed, some people may develop unusual levels of some body salts in the blood, such as potassium and uric acid, and this may lead to changes in kidney function, abnormal heartbeat or seizures.

The risk for TLS is in the first days or weeks of treatment with this medicine as you increase your dose.

Your healthcare team will tell you more about TLS, and you can refer to this guide for reminders of when and where to get blood tests to check how you're doing on the treatment and for any signs of TLS (see the **Your Treatment Plan** section).

What are the symptoms of TLS?

Temporarily stop your treatment and immediately tell your healthcare team or pharmacist if you notice any of the following symptoms:



Only restart taking your medicine when instructed to do so by your healthcare team.

What will my healthcare team do to help minimise the possibility of TLS?

Before starting your treatment



Your healthcare team will have **assessed your risk** of developing TLS and produced a **plan to monitor** you accordingly.



Your healthcare team may also give you **medicines** to help prevent the build-up of uric acid in your body, a potential consequence of TLS.



The first dose

As a precaution, the **first dose may be given in hospital** for monitoring purposes, due to the potential risk of developing TLS. Your healthcare team may decide to monitor you in hospital on the first day of subsequent dose increases too.

During your treatment



If your healthcare team determines you are at risk of TLS you may **receive your treatment in hospital** so that, if required, you can be given fluids, have blood tests, and be monitored for possible side effects.



Your healthcare team will tell you **when you need to have your blood tested** to check for initial signs of TLS after you start taking this medicine.

What can I do to help minimise the risk of TLS?



Turn up for blood tests

It is important for you to keep your scheduled appointments for blood tests to check for signs of TLS, because the changes in your blood that could lead to TLS may have no noticeable symptoms.



Drink plenty of water

It is very important that you start drinking plenty of water 2 days before you begin taking your medicine and keep doing so throughout your treatment – at least 1.5 to 2 litres (L) per day.



Why drink 1.5 to 2 litres of water per day?

Drinking 1.5 to 2 litres of water per day will help to continuously remove cancer cell breakdown products from your blood, through your urine.

What counts towards the 1.5 to 2 litres of water a day?

You can include other non-alcoholic and noncaffeinated drinks in this amount, but you must not drink grapefruit, Seville orange (bitter orange) or star fruit (carambola) juices.

Do I have to keep drinking at least 1.5 to 2 litres of water a day?

Yes, you should continue to drink the same amount each day while your doctor gradually increases the dose to the full standard dose, and beyond.



Your treatment plan

Keeping track over the first 5 weeks

The treatment plan template on the following pages has been designed to help you keep track of when and where to take your tablets, whether you need to have blood tests done, and other special instructions that your healthcare team has for you over those 5 weeks.

we	ek 1				This week's		
Day 0	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	
ate	Date	Date	Date	Date	Date	Date	Date
ttend	Take mg	Attend	Take mg	Take mg	Take mg	Take mg	Take
: (time)	at (time)	at (time)	at (time)	at (time)	at (time)	at (time)	at (time)
or blood test and review (await confirmation to	Attend	Do not take VENCLYXTO® until blood test	24 hours after the last dose	24 hours after the last do			
oceed the next morning)	at (time) For your 6-8 hour	review is complete Take mg					Attend at (time)
	blood test	at (time)					For blood test and revie (await confirmation to proceed with the new VENCLYXTO® dose)
Notes							
Drink at least 1.5 to 2 litres of water every day as directed by your doctor (tick off each glass as you go)	9999 9999	aaaa aaaa	9999 9999	9999 9999	a a a a a a a a	AAAA AAAA	9999 9999

Using your treatment plan

There is a page for each week of the first 5 weeks of treatment. With your healthcare team, fill in the tables for each week so you know exactly what to expect as you begin your treatment. You can also tick off each day to keep track of when you have taken your tablets.

Your plan will be tailored to your individual needs.

- Keep this guide somewhere visible so you have easy access to your treatment plan.
- Remember your progress by ticking off each step as you complete it.
- Take this guide with you when you return to the healthcare centre.



Please **read the information booklet** that explains your medication and treatment routine before using this treatment plan. Please refer to your **Package Leaflet** (**PL**) for additional information and your healthcare team if you have questions.



Day 0	Day 1	Day 2	Day 3
Date	Date	Date	Date
Attend	Take	Attend	Take
at (time)	at (time)	at (time)	at (time)
For blood test and review (await confirmation to proceed the next morning)	Attend at (time)	Do not take VENCLYXTO® until blood test review is complete	24 hours after the last dose
	at (time)	Take	
	For your 6-8 hour blood test	mg at (time)	
Notes			
	Medication taken	Medication taken	Medication taken
Uric acid reducing medicine (if prescribed)?	Yes 🖌 No 🖌	Yes 🖌 No 🖌	Yes 🖌 No 🖌

This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
Medication taken 🗸	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	
Yes 🖌 No 🗸	Yes 🖌 No 🏑	Yes 🖌 No 🖌	Yes 🖌 No 🖌	

Day 0	Day 1	Day 2	Day 3
Date	Date	Date	Date
Attend	Take mg	Attend	Take
at (time)	at (time)	at (time)	at (time)
For blood test and review (await confirmation to proceed the next morning)	Attend	Do not take VENCLYXTO® until blood test review is complete	24 hours after the last dose
	at (time)	Take	
	For your 6-8 hour blood test	mg at (time)	
Notes			
	Medication taken 📈 Dose taken 🛛 mg	Medication taken 📈 Dose taken 🛛 mg	Medication taken
Uric acid reducing medicine (if prescribed)?	Yes 🖌 No 🗸	Yes 📈 No 🗸	Yes 🖌 No √

This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
Medication taken 🗸	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	
Yes 🖌 No 🗸	Yes 🖌 No 🏑	Yes 🖌 No 🖌	Yes 🖌 No 🖌	



This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg	Take mg	Take mg	Take mg	
at (time)	at (time)	at (time)	at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose Attend at (time) For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
TTTT TTTT	YYY YYY	YYY YYY	TTTT TTTT	
Medication taken 📈 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken	
Yes 🗸 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🗸 No 🗸	



This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg	Take mg	Take mg	Take mg	
at (time)	at (time)	at (time)	at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose Attend at (time) For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
TTTT TTTT	YYY YYY	YYY YYY	TTTT TTTT	
Medication taken 📈 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken	
Yes 🗸 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🗸 No 🗸	



This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend at (time) For blood test and review	
			(await confirmation to proceed with the new VENCLYXTO® dose)	
U U U U U U U U	UUUU UUUU	UUUU UUUU	TTTT TTTT	
Medication taken Medication taken mg	Medication taken 📈 Dose taken mg	Medication taken 🗸	Medication taken Dose takenmg	
Yes 🖌 No 🖌	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🖌	

Day 0	Day 1	Day 2	Day 3
	Date	Date	Date
	Take	Take	Take
	mg at (time)	mg at (time)	mg at (time)
	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose
Notes			
Drink at least 1.5 to 2			
litres of water every day as directed by your doctor (tick off each			
glass as you go)			
Tick the box once you have taken your medicine, and make a	Medication taken	Medication taken	Medication taken
note of the dose taken	Dose taken mg	Dose taken mg	Dose taken mg
Uric acid reducing medicine (if prescribed)?	Yes 🖌 No 🖌	Yes No	Yes 🖌 No 🖌

This week's VENCLYXTO[®] dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend at (time) For blood test and review	
			(await confirmation to proceed with the new VENCLYXTO® dose)	
U U U U U U U U	UUUU UUUU	UUUU UUUU	TTTT TTTT	
Medication taken Medication taken mg	Medication taken 📈 Dose taken mg	Medication taken 🗸	Medication taken Dose takenmg	
Yes 🖌 No 🖌	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🖌	

Day 0	Day 1	Day 2	Day 3	
	Date	Date	Date	
	Take mg at (time)	Take mg at (time)	Take mg at (time)	
	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
Notes				
Drink at least 1.5 to 2 litres of water every day as directed by your doctor (tick off each glass as you go)	VVV VVVV	VVV VVVV	TTT TTTT	
Tick the box once you have taken your medicine, and make a note of the dose taken	Medication taken 📈 Dose taken mg	Medication taken 📈 Dose taken 🛛 mg	Medication taken	
Uric acid reducing medicine (if prescribed)?	Yes 🖌 No 🖌	Yes 🖌 No √	Yes 🖌 No 🏑	
Day 4	Day 5	Day 6	Day 7	
------------------------------	-------------------------------------	-------------------------------------	---	--
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend	
			at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO [®] dose)	
99999 99999	YYY YYY YYY	YYY YYY	YYY YYY	
Medication taken 🗸	Medication taken 📈 Dose taken mg	Medication taken 📈 Dose taken mg	Medication taken 🗸	
Yes 🖌 No 🗸	Yes 📈 No 🗸	Yes 🗸 No 🗸	Yes 📈 No 🖌	



Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend	
			at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO [®] dose)	
99999 99999	YYY YYY YYY	YYY YYY	YYY YYY	
Medication taken 🗸	Medication taken 📈 Dose taken mg	Medication taken 📈 Dose taken mg	Medication taken 🗸	
Yes 🖌 No 🗸	Yes 📈 No 🗸	Yes 🗸 No 🗸	Yes 📈 No 🖌	

Week 5



Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg	Take	Take mg	Take mg	
at (time)	at (time)	at (time)	at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend	
			at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
UUUU UUUU	UUUU UUUU	U U U U U U U U U U	U U U U U U U U U U	
Medication taken 🗸	Medication taken 📈 Dose taken mg	Medication taken 🖌 Dose taken mg	Medication taken 🖌 Dose taken mg	
Yes 🖌 No 🗸	Yes 🗸 No 🗸	Yes 🖌 No 🗸	Yes 🗸 No 🗸	

Week

Day 0	Day 1	Day 2	Day 3
	Date	Date	Date
	Take	Take	Take
	mg at (time)	mg at (time)	mg at (time)
	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose
Notes			
Drink at least 1.5 to 2 litres of water every day as directed by your doctor (tick off each glass as you go)	YYY YYY	YYY YYY	U U U U U U U U
Tick the box once you have taken your medicine, and make a note of the dose taken	Medication taken 🗸	Medication taken 🗸 Dose taken mg	Medication taken
Uric acid reducing medicine (if prescribed)?	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🖌

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg	Take mg	Take mg	Take mg	
at (time)	at (time)	at (time)	at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend	
			at (time)	
			For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
UUUU UUUU	UUUU UUUU	UUUU UUUU	U U U U U U U U U U	
Medication taken 🗸	Medication taken 📈 Dose taken mg	Medication taken 📈 Dose taken mg	Medication taken 🖌 Dose taken mg	
Yes 🖌 No 🗸	Yes 🗸 No 🗸	Yes 🖌 No 🗸	Yes 🗸 No 🗸	



Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose Attend at (time) For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
TTTT TTTT	YYY YYY	YYY YYY	TTT TTTT	
Medication taken 📈 Dose taken mg	Medication taken 📈 Dose taken 🛛 mg	Medication taken 📈 Dose taken mg	Medication taken	
Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 🖌 No 🗸	Yes 📝 No 🧹	



This week's VENCLYXTO® dose

Day 4	Day 5	Day 6	Day 7	
Date	Date	Date	Date	
Take mg at (time)	Take mg at (time)	Take mg at (time)	Take mg at (time)	
24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	24 hours after the last dose	
			Attend at (time) For blood test and review (await confirmation to proceed with the new VENCLYXTO® dose)	
TTTT TTTT	YYY YYY	YYY YYY	TTT TTTT	
Medication taken 🖌 Dose taken mg	Medication taken	Medication taken	Medication taken	
Yes 🗸 No 🖌	Yes 🖌 No 🗸	Yes 📈 No 🖌	Yes 🖌 No 🖌	

Treatment plan, week 6 and beyond

Once you've reached the 'maintenance dose' of 400mg per day, VENCLYXTO[®] can usually be taken at home on an ongoing basis.

Just remember to take it at around the same time each day, and refer to the important dosing notes on pages 4 to 9 whenever you need a reminder of how and when to take your medicine.



400mg each day (4x100mg tablets)



Remember to tell your healthcare team about any changes in your health, including

- Any signs of an upper respiratory tract infection, such as a runny nose, sore throat, cough or chest infection
- Feeling sick or being sick (nausea or vomiting)
- Feeling unusually tired
- Pneumonia
- Urinary tract infection

- Diarrhoea
- Constipation

• Fever or chills

At the first sign of infection or feeling unwell, contact your healthcare team.





Important reminder: If you start to take any other new medicine at any point whilst you are taking your medicine, consult your healthcare team.

Notes

46



Your patient information booklet

Useful information for healthcare professionals and services

On the opposite page you will find a credit card-sized booklet. Please detach this little booklet and carry it with you. It may fit in your purse or wallet.

This booklet contains useful information about VENCLYXTO® for healthcare professionals who usually work in other fields of medicine but are involved in your care. Show it to anyone who provides you with healthcare services such as pharmacists, dentists, opticians, practice nurses at your GP's clinic, paramedics etc.

It explains what this medicine is, the contraindications for treatment, potential drug interactions, and warnings and precautions to take.



VENCLYXTO® (venetoclax) Patient Booklet

This booklet needs to be shown in all interactions with healthcare professionals

I am currently receiving treatment with VENCLYXTO[®] for chronic lymphocytic leukaemia (CLL). It is a once-daily oral medication.¹

VENCLYXTO® may be given in combination with other medicines or alone.²



Reference VENCLYXTO® Package Leaflet available at www.medicines.ie

VENCLYXTO® is a registered trademark of AbbVie This booklet has been funded and produced by AbbVie Ltd Date of preparation: November 2023 | IE-VNCCLL-230033 © 2023 AbbVie Ltd

